

CanTEST have tested a total of **4864** samples to 20 June 2025. This month we provided **245** health & AOD brief interventions and **5/149** samples were voluntarily discarded by people after they received their results.

Benzos

Alprazolam (or Xanax/Mylan*)

was expected in **9** samples and detected in **3**.

1 sample also contained **bromazolam**, **diazepam** and **protonitazene**.

RED
Community Notice
3 June

1 sample also contained **4F-MBZP**, **AB-MDMSBA** and **N-cyclohexyl pentylone**.

YELLOW
Community Notice
3 July

1 sample also contained **bromazolam** and **flubromazepam**.

YELLOW
Community Notice
23 July

Where **alprazolam** was expected but not detected:

1 sample contained **clonazolam**, **bromazolam** and **bromonordiazepam**,

3 samples contained **clonazolam** and **bromazolam** and
2 samples contained **bromazolam** instead.

Diazepam (or Valium*)

was expected in **4** samples and detected in **3**.

1 sample also contained **nitrazepam** and **tapentadol**.

Where **diazepam** was expected but not detected,
1 sample contained **ethylbromazolam** instead.

Flunitrazepam

was expected and detected in **1** sample.

In **2** unknown benzo samples, **1** contained **diazepam** and **nitrazepam** and **1** contained **bromazolam**.

(The number of novel benzos identifiable onsite at CanTEST has increased recently, as our methods and libraries have improved)

*In some cases where service users provided an expected pharmaceutical brand name instead of an expected active drug.

Ketamine

was expected in **18** samples and detected in **13**.

Purities ranged from **22 - 86%**

1 sample also contained **procaine** and **tiletamine**.

Where **ketamine** was not detected, **1** sample contained **MDMA**, **procaine** and **tiletamine**, **1** sample contained **N-cyclohexyl benzylamine**, **procaine** and **tiletamine**, **2** samples contained **procaine** and **tiletamine**, and **1** contained **MDMA**.

MDMA

was expected in **35** samples and detected in **34**.

Samples tested and their purity ranges included:

10/10 capsules : **20 - 83%**

19/20 crystals : **52 - 84%**

4/4 pills : **16 - 40%**

1/1 powder : **56%**

1 sample also contained **paracetamol**.

Where **MDMA** was expected but not detected, **1** sample contained **ketamine** instead.

Heroin was expected and detected in **6** samples.

Purities ranged from **54 - 87%**

In addition to **heroin**, **3** samples also contained **6-MAM** and **acetylcodeine**.

Amphetamine was expected in **1** sample where **caffeine** was detected instead.

Cocaine was expected and detected in **32** samples. Purities ranged from **30 - 89%**
In addition to **cocaine**, **1** sample also contained **caffeine**.

LSD was expected in **4** samples and detected in **3**.

Methamphetamine was expected and detected in **15** samples. Purities ranged from **46 - 80%**.

2C-B was expected and detected in **3** samples. Purities ranged from **<5 - 79%**
In addition to **2C-B**, **1** sample also contained **MDMA**.

Tapentadol was expected and detected in **1** sample.

RED
Community Notice
31 May

Oxycodone was expected in **1** sample where **protonitazene** was detected instead.

Mephedrone was expected in **1** sample where **dimethylpentylone** was detected instead.

GHB/GBL/1,4-butanediol was expected in **1** sample where **1,4-butanediol** was detected.

DMXE was expected and detected in **1** sample.

Pregabalin was expected and detected in **1** sample.